Histoacryl®

The real thing Tissue adhesive



Biosurgicals



Histoacryl®

The real thing - now even simpler and faster

For decades medical professionals around the world have put their trust in Histoacryl*, the first surgical tissue adhesive based on cyanoacrylate. The successful application of Histoacryl* is described in more than 1,200 publications.

Now the classic product for the closure of skin wounds is available

- in a new ampoule with a twist-off tip.
- with increased content of n-butyl-2cyanoacrylate.
- with the indication for closure of skin wounds and for sclerosation therapy of esophageal and fundal varices.

New twist-off tip

Histoacryl®

now even simpler and faster

Simply twist off the ribbed tip of the ampoule

 Then, as usual, Histoacryl[®] can be applied precisely with the slim cannula



Translucent Histoacryl[®] L

for facial applications

To be on the safe side, it is recommended to use Histoacryl[®] L, a version which does not contain dye, for facial applications.



www.histoacryl.com

Classic advantages

Well-known advantages of Histoacryl[®]:

- fast wound closure
- superior strength
- simple and precise dosage due to slim cannula and blue dye
- applying just one layer is enough
- choice between blue and translucent version
- showering possible



Why tissue adhesive?

- no suturing no pain
- saves time and costs (no local anaesthetics, no suture removal, no second consultation)
- antibacterial film protects the wound
- no wound dressing necessary



Product Range

Ordering Information

Description	Color	Article No.
5 x 0.5 ml Histoacryl*	blue	105 0052
10 x 0.5 ml Histoacryl*	blue	105 0044
5 x 0.2 ml Histoacryl"	blue	105 0036
10 x 0.2 ml Histoacryl®	blue	105 0028
5 x 0.5 ml Histoacryl* L	translucent	105 0060

Literature is available upon request.

Wound closure in a minute:



Clean the wound and turn open the ampoule



Apply Histoacryl[®] sparingly, do not allow tissue adhesive to flow into the wound

Keep wound edges aligned for about one minute – that's all!

Histoacryl®



Adapt wound edges (wounds under tension and wounds longer than 3 cm require additional suturing)



Histoacryl®

The real thing - now even simpler and faster

Histoacryl[®] for closure of endoscopic incisions

The Histoacryl^{*} ampoule is supplied in sterile condition; therefore, Histoacryl^{*} is the ideal tissue adhesive for use in the OR.

For example, it may be used for the closure of endoscopic incisions, as described in:

Rosin D et al. Closure of laparoscopic trocar site wounds with cyanoacrylate tissue glue: a simple technical solution. J Laparoendosc Adv Surg Tech 2001; 11(3):157–9.



Sclerosation of esophageal and fundal varices with Histoacryl®

The formation of esophageal and fundal varices is a common and dangerous consequence of portal hypertension. Effective sclerosation of these varices is possible with Histoacryl[®].

Varically injected Histoacryl[®] polymerizes intravasally to form a plastic cylinder and effects immediate obturation or thrombosing of the vessel. The procedure is described in detail in the relevant literature, for instance in the following article:

Caldwell SH et al. Enbucrilate for gastric varices: extended experience in 92 patients. Aliment Pharmacol Ther 2007; 26:49–59.



Histoacryl®



B. Braun Surgical SA | Carretera de Terrassa, 121 | 08191 Rubí | Spain Phone +34 93 5 86 62 00 | Fax +34 93 6 99 73 03 | www.bbraun.es

Aesculap AG | Am Aesculap-Platz | 78532 Tuttlingen | Germany Phone +49 7461 95-0 | Fax +49 7461 95-26 00 | www.aesculap.com

Aesculap - a B. Braun company

Subject to technical changes. All rights reserved. This brochure may only be used for the exclusive purpose of obtaining information about our products. Reproduction in any form partial or otherwise is not permitted.

Brochure No. B01102